

Exercise Set 3

Exercise 3.1. A set of students applies for a set of seminars. Each student chooses exactly three seminars. Two seminars are chosen by 40 students, all others by fewer.

- (a) Prove that each student can be assigned to a seminar they chose without assigning more than 13 students to any seminar.
- (b) Show how to compute such an assignment in $\mathcal{O}(n^2)$ time, where n is the number of seminars.

(3+1 points)

Exercise 3.2. Let G be a graph and M a matching in G that is not maximum. In this exercise we use the terminology *disjoint subgraphs/paths/circuits* and mean it quite literally: Two subgraphs are *disjoint* if they have no edges and no vertices in common. (Note that the term *vertex-disjoint paths* is often used to mean that two paths have no *inner*-vertices in common, but possibly endpoints.)

- (i) Show that there are $\nu(G) - |M|$ disjoint M -augmenting paths in G .
- (ii) Show the existence of an M -augmenting path of length at most $\frac{\nu(G)+|M|}{\nu(G)-|M|}$.
- (iii) Let P be a shortest M -augmenting path in G and P' an $(M \Delta E(P))$ -augmenting path. Prove $|E(P')| \geq |E(P)| + 2 \cdot |E(P) \cap E(P')|$.

Consider the following algorithm: We start with the empty matching and in each iteration augment the matching along a shortest augmenting path. Let P_1, P_2, \dots be the sequence of augmenting paths chosen.

- (iv) Show that if $|E(P_i)| = |E(P_j)|$ for $i \neq j$, then P_i and P_j are disjoint.
- (v) Show that the sequence $|E(P_1)|, |E(P_2)|, \dots$ contains less than $2\sqrt{\nu(G)} + 1$ different numbers.

From now on, let G be bipartite and set $n := |V(G)|$ and $m := |E(G)|$.

- (vi) Given a non-maximum matching M in G show that we can find in $O(n+m)$ -time a family \mathcal{P} of disjoint shortest M -augmenting paths such that if M' is the matching obtained by augmenting M over every path in \mathcal{P} , then

$$\begin{aligned} \min\{|E(P)| : P \text{ is an } M'\text{-augmenting path}\} \\ > \min\{|E(P)| : P \text{ is an } M\text{-augmenting path}\} \end{aligned}$$

- (vii) Describe an algorithm with runtime $O(\sqrt{n}(m+n))$ that solves the CARDINALITY MATCHING PROBLEM in bipartite graphs.

(1+1+2+2+2+3+1=12 points)

Deadline: November 3, before the lecture. The websites for lecture and exercises can be found at:

https://ecampus.uni-bonn.de/goto_ecampus_crs_2772883.html

In case of any questions feel free to contact me at armbruster@or.uni-bonn.de.